

Supplementary Table 1. Peripheral blood analysis of *Nr4a3*^{-/-}*Nr4a1*^{-/-} mice at 2–3 weeks of age.

	Leukocytes (per μL) ($\times 10^3$) ¹	Erythrocytes (per μL) ($\times 10^6$) ¹	Hemoglobin (dL) ¹	Hematocrit (%) ¹	Platelets (per μL) ($\times 10^3$) ¹	Neutrophils	Lymphocytes	Monocytes (%) ²	Eosinophils	Immature/ Blast Forms
<i>Nr4a3</i> ^{+/+} <i>Nr4a1</i> ^{+/+} (<i>n</i> = 10) ³	2.3 \pm 0.4	6.7 \pm 0.1	12.6 \pm 0.2	39.4 \pm 0.3	794.3 \pm 50.2	22.9 \pm 1.4	69.1 \pm 1.6	0.9 \pm 0.2	4.7 \pm 0.5	2.5 \pm 0.34
<i>Nr4a3</i> ^{-/-} <i>Nr4a1</i> ^{+/-4}	3.2	5.2	10.1	33.5	550	36.8	39.1	0.7	13.7	9.8
	1.4	5.5	10.1	32.5	462	24.8	36.9	0.3	15.0	22.9
	1.6	3.7	6.7	22.2	621	19.6	64.7	0.3	5.2	10.5
	0.6	3.4	6.0	21.1	166	18.6	23.8	1.6	9.1	46.9
	2.2	2.2	4.3	14.2	493	30.2	27.3	1.0	4.8	36.7
	2.1	2.6	4.9	16.4	150	31.6	49.8	0	1.3	17.3
	1.9	3.8	6.7	23.4	77	15.4	60.3	0.3	0.3	20.8
	1.6	4.1	7.2	24.0	217	4.6	11.9	0	3.3	80.1
	2.8	3.4	6.6	21.5	305	19.4	68.3	0.7	6.8	4.5
	4.5	4.0	7.4	25.2	176	24.7	51.9	0.3	5.9	17.3
	5.8	4.2	8.1	27.6	403	18.0	42.7	0.3	6.7	32.6

¹ Values were determined by the Technicon Hematology System. ² Percent of cells was determined by identifying at least 300 leukocytes per blood smear. ³ Values are given as the mean \pm s.e.m. ⁴ Values for 11 individual mice are listed.